

7. Hrabětice

- 1CJ – cutting once per year – 15.6.
- 1CA – cutting once per year – 15.8.
- 2C - cutting twice per year – 15.6., 15.8.
- 2CAF - cutting twice per year – 15.6., 15.8. + wood ash fertilization (400 g/4m²)
- 2C2AF - cutting twice per year – 15.6., 15.8. + wood ash fertilization (double dose - 800 g/4m²)
- U - unmanaged treatment
- 2CM – cutting twice per year + manure
- 2CML – cutting twice per year + manure + liming
- Experiment was established in 2017
- Each plot has size 2 x 2 m
- Overall 4 randomized blocks (20 plots)
- *Trisetion* meadow
- Dominant species: *Festuca rubra*, *Agrostis capillaris*, *Bistorta major*, *Cirsium heterophyllum* and *Hypericum maculatum*

Plan of the experiment

	1CJ	1CA	2C	2CF	U	2C2F	2CM	2CML	SAMPLING
1. op.	1CJ	1CA	2C	2CF	U	2C2F	2CM	2CML	MONITORING
	2CF	2CM	1CJ	2CML	1CA	U	2C	2C2F	SAMPLING
2. op.	2CF	2CM	1CJ	2CML	1CA	U	2C	2C2F	MONITORING
	2C2F	1CA	U	2CM	2C	2CML	2CF	1CJ	SAMPLING
3. op.	2C2F	1CA	U	2CM	2C	2CML	2CF	1CJ	MONITORING
	2C	2CML	2C2F	1CJ	2CF	1CA	U	2CM	SAMPLING
4. op.	2C	2CML	2C2F	1CJ	2CF	1CA	U	2CM	MONITORING



